arXiv:1207.0466v1 [math.RA] 2 Jul 2012

Strongly J-$*$-Clean Rings

Huanyin Chen[*], Abdullah Harmancı[†], A. Çiğdem Özcan[‡]

Abstract

A $*$-ring R is strongly J-$*$-clean provided that for any $a \in R$, there exists a projection $e \in R$ such that $a - e \in J(R)$ and $ae = ea$. We prove, in this article, that a $*$-ring R is strongly J-$*$-clean, if and only if R is strongly $*$-clean ring and $R/J(R)$ is Boolean, if and only if R is uniquely clean and for any $a \in R$, $a - a^* \in J(R)$, if and only if for any $a \in R$, there exists a unique projection $e \in R$ such that $a - e \in U(R)$ and $ae = ea$, if and only if for any $a \in R$, there exists a unique projection $e \in R$ such that $a - e \in J(R)$. For any $*$-ideal I of R, it is shown that R is strongly J-$*$-clean if and only if R is abelian, every idempotent lifts modulo I and R/I is strongly J-$*$-clean. As consequences, many characterizations of such $*$-rings are obtained.

2010 Mathematics Subject Classification : 16W10, 16E50

Key words: strongly J-$*$-clean ring, strongly $*$-clean ring, uniquely clean ring.

1 Introduction

Let R be an associative ring with identity. A ring R is *strongly clean* provided that for any $a \in R$, there exists an idempotent $e \in R$ such that $a - e \in U(R)$ and $ae = ea$ ([2, 3, 11]). A ring R is *strongly J-clean* provided that for any $a \in R$, there exists an idempotent $e \in R$ such that $a - e \in J(R)$ and $ae = ea$ ([4]). A ring R is *uniquely clean* provided that for any $a \in R$, there exists a unique idempotent $e \in R$ such that $a - e \in U(R)$ ([5, 7, 12, 16]). A ring R is *uniquely strongly clean* provided that for any $a \in R$, there exists a unique idempotent $e \in R$ such that $a - e \in U(R)$ and $ae = ea$ ([8, 15, 17]). As is well known, {uniquely clean rings} $\subsetneqq$ {uniquely strongly clean rings} $\subsetneqq$ {strongly J-clean rings} $\subsetneqq$ {strongly clean rings}. For more references on strongly cleanness, we refer the reader to [6, 11].

An involution of a ring R is just an anti-automorphism whose square is the identity map 1_R. Thus an *involution* of a ring R is an operation $* : R \to R$ such that $(x + y)^* = x^* + y^*$, $(xy)^* = y^*x^*$ and $(x^*)^* = x$ for all $x, y \in R$. A ring R with involution $*$ is called a *$*$-ring*. An element e in a $*$-ring R is called a *projection* if $e^2 = e = e^*$ (cf. [1]). Parallel to the preceding strongly clean rings, strongly $*$-clean rings are studied recently. A $*$-ring R is *strongly $*$-clean* if each of its elements is the sum of a unit and a projection that commute with each other

---

[*] Department of Mathematics, Hangzhou Normal University, Hangzhou, 310036, People's Republic of China, e-mail: huanyinchen@yahoo.cn

[†] Hacettepe University, Department of Mathematics, 06800 Beytepe Ankara, Turkey, e-mail: abdullahharmanci@gmail.com

[‡] Corresponding author. Hacettepe University, Department of Mathematics, 06800 Beytepe Ankara, Turkey, e-mail: ozcan@hacettepe.edu.tr

([10, 14]). Our motivation is to consider uniquely strongly cleanness and strongly J-cleanness for $*$-rings. Surprisingly, we will observe that these ones coincide with each other for $*$-rings.

We say that a $*$-ring is *strongly J-$*$-clean* provided that for any $a \in R$, there exists a projection $e \in R$ such that $a - e \in J(R)$ and $ae = ea$. We will investigate the behaving of involutions and get the relations among strongly $*$-cleanness, uniquely cleanness and strongly J-$*$-cleanness. Here it is proved that a $*$-ring R is strongly J-$*$-clean, if and only if R is strongly $*$-clean ring and $R/J(R)$ is Boolean, if and only if R is uniquely clean and for any $a \in R$, $a - a^* \in J(R)$, if and only if for any $a \in R$, there exists a unique projection $e \in R$ such that $a - e \in U(R)$ and $ae = ea$, if and only if for any $a \in R$, there exists a unique projection $e \in R$ such that $a - e \in J(R)$. For any $*$-ideal I of R which is contained in $J(R)$, it is shown that R is strongly J-$*$-clean if and only if R is abelian, every idempotent lifts modulo I and R/I is strongly J-$*$-clean. As consequences, various structure properties and examples of such new class of $*$-rings are obtained.

Throughout, all rings are associative with identity. We use $U(R)$ to denote the set of all invertible elements in R. $J(R)$ stands for the Jacobson radical of R.

2 Strongly $*$-Clean Rings

A $*$-ring R is said to be *J-$*$-clean* provided that for any $a \in R$ there exists a projection $e \in R$ such that $a - e \in J(R)$. The purpose of this section is to characterize strongly J-$*$-clean rings by means of strongly $*$-cleanness. We begin with the following elementary result.

Proposition 2.1 *Let R be a $*$-ring. Then the following are equivalent:*

(1) *R is strongly J-$*$-clean.*

(2) *R is abelian and R is J-$*$-clean.*

(3) *R is strongly J-clean and strongly $*$-clean.*

Proof (1) $\Rightarrow$ (3) Clearly, R is strongly J-clean. For any $a \in R$, there exists a projection $e \in R$ such that $u := a - e \in J(R)$ and $ae = ea$. Then $a = (1 - e) + (2e - 1 + u)$. Clearly, $1 - e \in R$ is a projection. As $(2e - 1)^2 = 1$, we see that $2e - 1 + u \in U(R)$. Thus, R is strongly $*$-clean.

(3) $\Rightarrow$ (2) Since R is strongly $*$-clean, it follows from [10, Theorem 2.2] that R is abelian and every idempotent is a projection. Therefore R is J-$*$-clean.

(2) $\Rightarrow$ (1) is obvious. □

Example 2.2 (1) Let the ring $R = \mathbb{Z}_2 \oplus \mathbb{Z}_2$. Define $\sigma : R \to R$ by $\sigma(x, y) = (y, x)$. Consider the ring $T_2(R, \sigma) = \left\{ \begin{pmatrix} a & b \\ 0 & a \end{pmatrix} \mid a, b \in R \right\}$ with the following operations:

$$\begin{pmatrix} a & b \\ 0 & a \end{pmatrix} + \begin{pmatrix} c & d \\ 0 & c \end{pmatrix} = \begin{pmatrix} a+c & b+d \\ 0 & a+c \end{pmatrix}, \quad \begin{pmatrix} a & b \\ 0 & a \end{pmatrix} \cdot \begin{pmatrix} c & d \\ 0 & c \end{pmatrix} = \begin{pmatrix} ac & ad + b\sigma(c) \\ 0 & ac \end{pmatrix}.$$

Then $J(T_2(R,\sigma)) = \{ \begin{pmatrix} 0 & b \\ 0 & 0 \end{pmatrix} \mid b \in R\}$ and it is nilpotent. In addition, $T_2(R,\sigma)/J(T_2(R,\sigma)) \cong R$ is Boolean. Define $* : R \to R$ by $\begin{pmatrix} a & b \\ 0 & a \end{pmatrix}^* = \begin{pmatrix} a & \sigma(b) \\ 0 & a \end{pmatrix}$. As $\sigma^2 = 1_R$, we check easily that $*$ is an involution of $T_2(R,\sigma)$. Since, for any $a, b \in R$, $\begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix}$ is a projection and $\begin{pmatrix} a & b \\ 0 & a \end{pmatrix} - \begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix} \in J(T_2(R,\sigma))$, $T_2(R,\sigma)$ is a J-$*$-clean ring but it is not strongly J-$*$-clean since $\begin{pmatrix} (0,1) & (0,0) \\ (0,0) & (0,1) \end{pmatrix}$ is not central.

(2) Let $R = \mathbb{Z}_{(3)}$ be the localization, and $* = 1_R$, the identical automorphism of R. Since R is local, it is strongly $*$-clean but not strongly J-$*$-clean because $R/J(R)$ is not Boolean, for example $(\frac{2}{1})^2 - \frac{2}{1} \notin J(R) = 3R$.

(3) Let $R = \begin{pmatrix} \mathbb{Z}_2 & \mathbb{Z}_2 \\ 0 & \mathbb{Z}_2 \end{pmatrix}$. Then R is a strongly J-clean ring, but R cannot be strongly J-$*$-clean for any involution $*$ on R.

(4) Let $R = \{ \begin{pmatrix} a & b \\ c & a \end{pmatrix} \mid a, b, c \in \mathbb{Z}_2\}$. Then R is a commutative ring with the following operations:

$$\begin{pmatrix} a & b \\ c & a \end{pmatrix} + \begin{pmatrix} a' & b' \\ c' & a' \end{pmatrix} = \begin{pmatrix} a+a' & b+b' \\ c+c' & a+a' \end{pmatrix},$$

$$\begin{pmatrix} a & b \\ c & a \end{pmatrix} \cdot \begin{pmatrix} a' & b' \\ c' & a' \end{pmatrix} = \begin{pmatrix} aa' & ab'+ba' \\ ca'+ac' & aa' \end{pmatrix}.$$

Define $* : R \to R$ by $\begin{pmatrix} a & b \\ c & a \end{pmatrix}^* = \begin{pmatrix} a & c \\ b & a \end{pmatrix}$. Then R is a $*$-ring and the projections of R are the zero and the identity matrices. Also $J(R) = \{ \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \}$. Hence R is a strongly J-$*$-clean ring.

Let R be a $*$-ring, and let $C(R) = \{a \in R \mid ax = xa \text{ for any } x \in R\}$. Then $C(R)$ is a subring of R. It is easy to check that $* : C(R) \to C(R)$ is also an anti-automorphism. Thus, $C(R)$ is a $*$-ring.

Corollary 2.3 *Let R be a $*$-ring. Then R is strongly J-$*$-clean if and only if*

(1) *$C(R)$ is J-$*$-clean;*

(2) *$R = C(R) + J(R)$.*

Proof Suppose that R is strongly J-$*$-clean. Then R is strongly $*$-clean by Proposition 2.1. Thus, every projection in R is central from [10, Theorem 2.2]. For any $a \in C(R)$, there exist a projection $e \in R$ and an element $u \in J(R)$ such that $a = e + u$. Hence, $e \in C(R)$ is a projection. This implies that $u \in J(R) \cap C(R) \subseteq J\big(C(R)\big)$. As a result, $C(R)$ is J-$*$-clean. On the other hand, it follows from Proposition 2.1 that R is abelian strongly J-clean. In view of [6, Corollary 16.4.16], R uniquely clean. According to [12, Proposition 25], $R = C(R) + J(R)$.

Conversely, assume that (1) and (2) hold. For any idempotent $e \in R$, there exist some $b \in C(R)$ and $c \in J(R)$ such that $e = b + c$. As $C(R)$ is J-$*$-clean, we can find a projection $f \in C(R)$ and a $w \in J\big(C(R)\big)$ such that $b = f + w$. Hence, $e = f + (w + c)$, and so $e - f = w + c$. Obviously, $(e - f)\big(1 - (e - f)^2\big) = 0$. On the other hand, $(e - f)^2 \in J\big(C(R)\big) + J(R)$, and so $1 - (e - f)^2 \in U(R)$. Therefore $e = f$, i.e., every idempotent in R is a projection. For any $a \in R$, write $a = s + t, s \in C(R), t \in J(R)$. We have a projection $g \in R$ such that $s = g + v$ with $v \in J\big(C(R)\big)$. This shows that $a = g + (t + v)$, and so R is J-$*$-clean. According to [10, Lemma 2.1] and Proposition 2.1, we complete the proof. $\square$

According to [5, Theorem 2.1], a ring R is uniquely clean if and only if R is an abelian exchange ring and $R/M \cong \mathbb{Z}_2$ for all maximal ideals M of R. We can derive an analogue for strongly J-$*$-clean rings.

Theorem 2.4 *Let R be a $*$-ring. Then R is strongly J-$*$-clean if and only if*

(1) *R is strongly $*$-clean ring;*

(2) *$R/M \cong \mathbb{Z}_2$ for all maximal ideals M of R.*

Proof Suppose that R is strongly J-$*$-clean. Then R is an abelian strongly J-clean ring by Proposition 2.1. Thus, R is uniquely clean from [6, Corollary 16.4.16]. Therefore $R/M \cong \mathbb{Z}_2$ for all maximal ideals M of R by [5, Theorem 2.1].

Conversely, assume that (1) and (2) hold. Then every idempotent in R is a projection. Further, R is an abelian exchange ring. According to [5, Theorem 2.1], R is uniquely clean, and so R is strongly J-clean. Therefore R is strongly J-$*$-clean by Proposition 2.1. $\square$

Corollary 2.5 *Let R be a $*$-ring. Then R is strongly J-$*$-clean if and only if*

(1) *R is strongly $*$-clean ring;*

(2) *For all maximal ideal M of R, 1 is not the sum of two units in R/M.*

Proof One direction is obvious. Conversely, assume that (1) and (2) hold. Let M be a maximal ideal of R. Clearly, R/M is an abelian exchange ring; hence, R/M is an exchange ring with artinian primitive factors. As R/M is simple, $J(R/M) = 0$. Let $f \in R/M$ be an idempotent. Then $(R/M)f(R/M) = 0$ or R/M, and so $f = \overline{0}$ or $\overline{1}$. This means that R/M is indecomposable. Therefore R/M is simple artinian. Thus, $R/M \cong M_n(D)$, where D is a division ring. As every idempotent in R/M is central, we deduce that $n = 1$, and so $R/M \cong D$. If $|D| \geq 3$, then we can find a set $\{0, 1, x\} \subseteq D$, where $x \neq 0, 1$. This shows that $1 - x \in U(R/M)$, a contradiction. Therefore $R/M \cong \mathbb{Z}_2$. According to Theorem 2.4, we complete the proof. $\square$

Theorem 2.6 *Let R be a $*$-ring. Then the following are equivalent:*

(1) *R is a strongly J-$*$-clean ring.*

(2) *$R/J(R)$ is Boolean and R is strongly $*$-clean.*

Proof (1) $\Rightarrow$ (2) In view of Proposition 2.1, R is strongly $*$-clean and R is strongly J-clean. By virtue of [6, Proposition 16.4.15], $R/J(R)$ is Boolean.

(2) $\Rightarrow$ (1) As R is strongly $*$-clean, it is strongly clean. According to [6, Proposition 16.4.15], R is strongly J-clean. In light of Proposition 2.1, R is a strongly J-$*$-clean ring. $\square$

Corollary 2.7 *Let R be a $*$-ring. Then R is a strongly J-$*$-clean ring if and only if*

(1) *R is a strongly $*$-clean ring;*

(2) *Every nonzero idempotent in R is not the sum of two units.*

Proof Suppose that R is a strongly J-$*$-clean ring. By virtue of Theorem 2.6, R is a strongly $*$-clean ring, and $R/J(R)$ is Boolean. Let $0 \neq e \in R$ be an idempotent. If $e = u + v$ for some $u, v \in U(R)$. Then $\overline{e} = \overline{u} + \overline{v}$ in $R/J(R)$. As $\overline{u} = \overline{v} = \overline{1}$, we see that $2 - e \in J(R)$. In light of [4, Proposition 3.1], $2 \in J(R)$; hence, $e \in J(R)$. This implies that $e = 0$, a contradiction. Therefore every nonzero idempotent in R is not the sum of two units.

Conversely, assume that (1) and (2) hold. Then R is an exchange ring. According to [9, Theorem 13], $R/J(R)$ is Boolean. Therefore we complete the proof from Theorem 2.6. $\square$

Recall that a ring R is *local* if R has only one maximal right ideal. As is well known, a ring R is local if and only if $a + b = 1$ in R implies that either a or b is invertible.

Corollary 2.8 *Let R be a local $*$-ring. Then the following are equivalent:*

(1) *R is strongly J-$*$-clean.*

(2) *R is strongly J-clean.*

(3) *R is uniquely clean.*

(4) *$R/J(R) \cong \mathbb{Z}_2$.*

(5) *1 is not the sum of two units in R.*

Proof Since R is a local $*$-ring, R is strongly $*$-clean. Therefore the result follows from Proposition 2.1, Theorem 2.4, Theorem 2.6 and [4, Lemma 4.2]. $\square$

A $*$-ring R is called *$*$-regular* if R is (von Neumann) regular and the involution is proper, equivalently for every x in R there exists a projection p such that $xR = pR$ (see [1]).

Corollary 2.9 *Let R be a $*$-regular ring. Then R is strongly J-$*$-clean if and only if R is Boolean.*

Proof Suppose that R is Boolean. Then $R/J(R)$ is Boolean. For any idempotent $e \in R$, there exists a projection p such that $eR = pR$. As R has stable range one, we have a unit $u \in R$ such that $e = pu$. Clearly, $u = 1$, and so $e = p$. This implies that R is strongly $*$-clean. According to Theorem 2.6, R is strongly J-$*$-clean.

Conversely, assume that R is strongly J-$*$-clean. Then $R/J(R)$ is Boolean by Theorem 2.6. But R is regular, and so $J(R) = 0$. Therefore R is Boolean. □

3 Structure Theorems

We can characterize strongly J-$*$-clean rings by means of uniquely cleanness as follows.

Theorem 3.1 *Let R be a $*$-ring. Then the following are equivalent:*

(1) *R is strongly J-$*$-clean.*

(2) *R is uniquely clean and for any $a \in R$, $a - a^* \in J(R)$.*

(3) *R is uniquely clean and for any $a \in R$, $a + a^* \in J(R)$.*

Proof (1) ⇒ (2) In view of Proposition 2.1, R is an abelian strongly J-clean ring, and so it is uniquely clean. For any $a \in R$, there exist a projection $e \in R$ and an element $u \in J(R)$ such that $a = e + u, ae = ea$. Thus, $a^* = e^* + u^*$. As $\big(J(R)\big)^* \subseteq J(R)$, we see that $a - a^* = (e - e^*) + (u - u^*) \in J(R)$.

(2) ⇒ (3) Since R is uniquely clean, it follows from [12, Lemma 18], $2 \in J(R)$. For any idempotent e in R, $e - e^* \in J$. Since $(e - e^*)^3 = e - e^*$ we have that $(e - e^*)(1 - (e - e^*)^2) = 0$. Hence, $e = e^*$, i.e. every idempotent is a projection. For any $a \in R$, there exist a projection $e \in R$ and an element $u \in J(R)$ such that $a = e + u$, $ae = ea$ (see [12, Theorem 20]). Thus, $a^* = e^* + u^*$. As a result, we deduce that $a + a^* = (e + e^*) + (u + u^*) = 2e + (u + u^*) \in J(R)$.

(3) ⇒ (1) In view of [6, Corollary 16.4.16], R is abelian strongly J-clean. For any idempotent $e \in R$, $e + e^* \in J(R)$. Thus, $(e - e^*)(e + e^*)^2 = e - e^*$; hence, $(e - e^*)\big(1 - (e + e^*)^2\big) = 0$. This implies that $e = e^*$, i.e., every idempotent is a projection. Hence, R is strongly J-$*$-clean. □

Let R be a $*$-ring. Then $*$ induces an involution of the power series ring $R[[x]]$, denoted by $*$, where $\big(\sum\limits_{i=0}^{\infty} a_i x^i\big)^* = \sum\limits_{i=0}^{\infty} a_i^* x^i$. Likewise, $*$ induces an involution of $R[[x]]/(x^n)$ $(n \geq 1)$, denoted by $*$, where $(\sum\limits_{i=0}^{n-1} a_i x^i)^* = \sum\limits_{i=0}^{n-1} a_i^* x^i$.

Corollary 3.2 *Let R be a $*$-ring. Then the following are equivalent:*

(1) *R is a strongly J-$*$-clean ring.*

(2) *$R[[x]]$ is a strongly J-$*$-clean ring.*

(3) *$R[[x]]/(x^2)$ is a strongly J-$*$-clean ring.*

(4) *$R[[x]]/(x^n)$ is a strongly J-$*$-clean ring for all $n \geq 2$.*

Proof (1) $\Rightarrow$ (2) Since R is strongly J-$*$-clean, $R[[x]]$ is strongly $*$-clean by [10, Corollary 2.10]. $R[[x]]$ is also strongly J-clean by [6, Example 16.4.17]. Hence $R[[x]]$ is strongly J-$*$-clean by Proposition 2.1.

(2) $\Rightarrow$ (4) In view of Theorem 3.1, $R[[x]]$ is uniquely clean, and then so is $R[[x]]/(x^n)$ by [12, Theorem 22]. Obviously, $S := R[[x]]/(x^n) = \{\sum\limits_{i=0}^{n-1} a_ix^i \mid a_0, \cdots, a_{n-1} \in R, x^n = 0\}$. For any $f \in S$, it follows from Theorem 3.1 that $f(0) - \big(f(0)\big)^* \in J(R)$, and so $f - f^* = f(0) - (f(0))^* + \sum\limits_{i=1}^{n-1} b_ix^i \in J(S)$. By using Theorem 3.1 again, $R[[x]]/(x^n)$ is a strongly J-$*$-clean ring.

(4) $\Rightarrow$ (3) is trivial.

(3) $\Rightarrow$ (1) Let $S = R[[x]]/(x^2)$. Then $S = \{a + bx \mid a, b \in R, x^2 = 0\}$. For any $a + bx \in S$, there exists a projection $e + fx \in S$ such that $(a + bx)(e + fx) = (e + fx)(a + bx)$ and $(a + bx) - (e + fx) \in J(S)$. This implies that $e \in R$ is a projection, $ae = ea$ and $a - e \in J(R)$. Therefore R is a strongly J-$*$-clean ring. $\square$

Let R be a $*$-ring. Then R is a strongly J-$*$-clean ring if and only if $R[x]/(x^2)$ is a strongly J-$*$-clean ring. Obviously, $R[x]/(x^2) \cong R[[x]]/(x^2)$, and we are done.

Theorem 3.3 *Let R be a $*$-ring. Then the following are equivalent:*

(1) *R is strongly J-$*$-clean.*

(2) *R is uniquely clean and R is strongly $*$-clean.*

(3) *For any $a \in R$, there exists a unique projection $e \in R$ such that $a - e \in U(R)$ and $ae = ea$.*

(4) *For any $a \in R$, there exists a unique projection $e \in R$ such that $a - e \in J(R)$.*

Proof (1) $\Rightarrow$ (2) is clear from Theorem 3.1 and Proposition 2.1.

(2) $\Rightarrow$ (3) For any $a \in R$, it follows from [12, Theorem 20] that there exists a unique idempotent $e \in R$ such that $a - e \in J(R)$. Since R is strongly $*$-clean, e is a central projection. Therefore there exists a unique projection $e \in R$ such that $a - e \in U(R)$ and $ae = ea$.

(3) $\Rightarrow$ (1) Clearly, R is strongly $*$-clean. In light of [10, Theorem 2.2], R is abelian and every idempotent in R is a projection. Thus, R is uniquely clean. Let $a \in R$. According to [12, Theorem 20], there exists an idempotent $e \in R$ such that $a - e \in J(R)$. Thus, we have a projection $e \in R$ such that $a - e \in J(R)$ and $ae = ea$. Therefore R is strongly J-$*$-clean.

(1) $\Rightarrow$ (4) is trivial.

(4) $\Rightarrow$ (1) For any idempotent e in R, there exists a projection $g \in R$ such that $e - g \in J(R)$. Thus, $e^* - g \in J(R)$. Therefore $e - e^* = (e - g) - (e^* - g) \in J(R)$. Clearly, there exists a projection $h \in R$ such that $2 = h + w$ where $w \in J(R)$. Thus, $1 - h = -1 + w \in U(R)$. Hence $1 - h = 1$, and so $h = 0$. This implies that $2 \in J(R)$. As a result, we deduce that $e + e^* = (e - e^*) + 2e^* \in J(R)$.

Set $z = 1 + (e - e^*)^*(e - e^*)$. Write $t = z^{-1}$. Since $z^* = z$, $t^* = t$. Also $e^*z = e^*ee^* = ze^*$, and so $e^*t = te^*$, and $et = te$. Set $f = e^*et = te^*e$. Then $f^* = f$, $f^2 = e^*ete^*et = e^*ee^*(tet) = e^*ztet = e^*et = f$, $fe = f$ and $ef = ee^*et = ezt = e$. Now $e = f + (e - f)$ and $e - f = e - e^*et = ee^*et - e^*et = (e - e^*)e^*et \in J(R)$. Here $f = f^* = f^2$. In addition, $f = e^*e[1 + (e^* - e)(e - e^*)]^{-1}$.

Set $z' = 1+(e^*-e)^*(e^*-e)$. Write $t' = (z')^{-1}$. Since $(z')^* = z'$, $(t')^* = t'$. Also $ez' = ee^*e = z'e$. Set $f' = ee^*t' = t'ee^*$. As in the preceding proof, we see that $f' = (f')^2 = (f')^*$ and $ef' = f'$, $f'e = e$. In addition, $e-f' = f'e-f' = t'ee^*(e-e^*) \in J(R)$ where $f' = [1+(e-e^*)(e^*-e)]^{-1}ee^*$.

Thus we get $e = f+(e-f) = f'+(e-f')$ with $e-f, e-f' \in J(R)$, f and f' are projections. By the uniqueness, we get $e^*e[1+(e^*-e)(e-e^*)]^{-1} = [1+(e-e^*)(e^*-e)]^{-1}ee^*$. This shows that $[1+(e-e^*)(e^*-e)]e^*e = ee^*[1+(e^*-e)(e-e^*)]$. Obviously, $(e-e^*)(e^*-e)e^*e = -e^*e+e^*ee^*e$ and $ee^*(e^*-e)(e-e^*) = -ee^*+ee^*ee^*$. Consequently, $e^*ee^*e = ee^*ee^*$. One easily checks that $(e-e^*)^3-(e-e^*) = -ee^*e+e^*ee^*$ and $[(e-e^*)^3-(e-e^*)](e+e^*) = (e-e^*)^3-(e-e^*)$. Thus $(e-e^*)((e-e^*)^2-1)((e+e^*)-1) = 0$. As $e-e^*$, $e+e^* \in J(R)$, we derive that $e = e^*$. Therefore every idempotent is a projection, and so R is abelian by [10, Lemma 2.1], and the result follows. □

We note that the unique projection in (3) or (4) of Theorem 3.3 can not be replaced by the unique idempotent even for a commutative $*$-ring as the following example shows:

Example 3.4 Let $R = \left\{ \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 0 & 1 \end{pmatrix} \right\}$ where $0, 1 \in \mathbb{Z}_2$. Define $* : R \to R$, $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \mapsto \begin{pmatrix} a+b & b \\ a+b+c+d & b+d \end{pmatrix}$. Then R is a commutative $*$-ring with the usual matrix addition and multiplication. In fact, R is Boolean, and so, for any $a \in R$, there exists a unique idempotent $e \in R$ such that $a-e \in U(R)$ (or, $a-e \in J(R)$) and $ae = ea$. But R is not strongly J-$*$-clean, even not a $*$-clean ring.

Theorem 3.5 *Let R be a $*$-ring. Then R is strongly J-$*$-clean if and only if*

(1) *R is strongly $*$-clean;*

(2) $J(R) = \{x \in R \mid 1-x \in U(R)\}$.

Proof Suppose that R is strongly J-$*$-clean. By virtue of Theorem 3.1, R is uniquely clean. Obviously, $J(R) \subseteq \{x \in R \mid 1-x \in U(R)\}$. Suppose that $1-x \in U(R)$. If $x \notin J(R)$, then $0 \neq xR \nsubseteq J(R)$. In view of [12, Lemma 17], there exists an idempotent $0 \neq e \in xR$. Write $e = xr$ for an $r \in R$. Then $e = (exe)(ere)$ as every idempotent in R is central. It is easy to see that R is directly finite. Thus, $exe \in U(eRe)$. In view of [12, Corollary 5], eRe is uniquely clean. Clearly, $0+exe = e+e(x-1)e$. The uniqueness implies that $0 = e$, a contradiction. Therefore $x \in J(R)$, and so $\{x \in R \mid x-1 \in U(R)\} \subseteq J(R)$. Thus, $J(R) = \{x \in R \mid 1-x \in U(R)\}$.

Conversely, assume that (1) and (2) hold. Let $a \in R$. Then we can find a projection $e \in R$ such that $(a-1)-e \in U(R)$ and $e(a-1) = (a-1)e$. That is, $(1-a)+e \in U(R)$. As $1-(a-e) \in U(R)$, by hypothesis, $a-e \in J(R)$. In addition, $ea = ae$. Therefore R is strongly J-$*$-clean. □

Corollary 3.6 *Let R be a $*$-ring. Then R is strongly J-$*$-clean if and only if*

(1) *R is strongly $*$-clean;*

(2) *For any $a \in R$, $a+a^* \in J(R)$;*

(3) $J(R) = \{x \in R \mid 1 + xx^* \in U(R)\}$.

Proof Suppose that R is strongly J-$*$-clean. By virtue of Proposition 2.1 and Theorem 3.1, R is a strongly $*$-clean ring with $2 \in J(R)$. For any $x \in R$, it follows from Theorem 3.1 that $x + x^* \in J(R)$. If $1 + xx^* \in U(R)$, then $(1 + x)(1 + x^*) = 1 + x + x^* + xx^* \in U(R)$. This implies that $1 + x \in R$ is right invertible. Similarly, $1 + x \in R$ is left invertible. In view of Theorem 3.5, $-x \in J(R)$, and so $x \in J(R)$. This shows that $J(R) = \{x \in R \mid 1 + xx^* \in U(R)\}$.

Conversely, assume that (1), (2) and (3) hold. If $1 + x \in U(R)$, then $1 + x^* \in U(R)$; hence, $(1 + x)(1 + x^*) \in U(R)$. As $x + x^* \in J(R)$, we see that $1 + xx^* \in U(R)$. This implies that $x \in J(R)$, and so $-x \in J(R)$. As a result, $J(R) = \{x \in R \mid 1 - x \in U(R)\}$. According to Theorem 3.5, the result follows. $\square$

Corollary 3.7 *Let R be a $*$-ring. Then R is strongly J-$*$-clean if and only if*

(1) *R is strongly $*$-clean;*

(2) *$2 \in J(R)$;*

(3) *$U\big(R/J(R)\big)$ is torsion.*

Proof If R is strongly J-$*$-clean, then R is strongly $*$-clean and $2 \in J(R)$. In addition, $R/J(R)$ is Boolean. Thus, $U\big(R/J(R)\big) = \{\overline{1}\}$ is torsion.

Conversely, assume that (1), (2) and (3) hold. Assume that $1 - x \in U(R)$. Then $\overline{1 - x} \in U\big(R/J(R)\big)$. By hypothesis, there exists some $n \in \mathbb{N}$ such that $(\overline{1 - x})^n = \overline{1}$, and so $(\overline{1 - x})^{2n} = \overline{1}$. As $2 \in J(R)$, we see that $x^{2n} \in J(R)$. Clearly, R is an abelian exchange ring, and then so is $R/J(R)$. In view of [13, Lemma 4.10], $R/J(R)$ is reduced, i.e. it has no nonzero nilpotent elements, and so $x \in J(R)$. This implies that $J(R) = \{x \in R \mid 1 - x \in U(R)\}$. According to Theorem 3.5, R is strongly J-$*$-clean. $\square$

Lemma 3.8 *Let R be a $*$-ring. Then R is strongly J-$*$-clean if and only if for any $a \in R$, there exists a unique idempotent $e \in R$ such that $a - e \in U(R), ae = ea, ae^* = e^*a$ and $e - e^* \in J(R)$.*

Proof Suppose that R is strongly J-$*$-clean. Then there exists a projection $f \in R$ such that $a - f \in U(R), af = fa, af^* = f^*a$ and $f - f^* = 0 \in J(R)$ by Theorem 3.3. Assume that there exists an idempotent $e \in R$ such that $u := a - e \in U(R), ae = ea, ae^* = e^*a$ and $e - e^* \in J(R)$. Let $p = 1 + (e^* - e)^*(e^* - e)$. As $ae = ea, ae^* = e^*a$, we see that $ap = pa$. Clearly, $p \in U(R)$. Write $q = p^{-1}$. Then $p^* = p$, and so $q^* = q$. Further, $ep = e(1 - e - e^* + ee^* + e^*e) = ee^*e = (1 - e - e^* + ee^* + e^*e)e = pe$. Thus, we see that $eq = qe$ and $e^*q = qe^*$. Set $g = ee^*q$. Then $g^2 = ee^*qee^*q = qee^*ee^*q = qpee^*q = ee^*q = g$. In addition, $g^* = q^*ee^* = ee^*q = g$, i.e., $g \in R$ is a projection. As $aq = qa$, we see that $ag = ga$. One easy checks that $eg = g$ and $ge = ee^*qe = ee^*eq = epq = e$. This implies that $e - g = e - ee^*p^{-1} = e\big(e(e^* - e)^* - 1\big)(e^* - e)p^{-1} \in J(R)$, and so $e - g + u \in U(R)$. By the uniqueness in Theorem 3.3, we see that $g = f$. This implies that $(e - f)^2 = (e - g)^2 = e - eg - ge + g = 0$. In view of Proposition 2.1, R is abelian, and so $(e - f)^3 = e - f$; hence, $e = f$. Therefore there exists a unique idempotent $e \in R$ such that $a - e \in U(R), ae = ea, ae^* = e^*a$ and $e - e^* \in J(R)$.

Conversely, let $a \in R$. Then there exists an idempotent $e \in R$ such that $u := a - e \in U(R), ae = ea, ae^* = e^*a$ and $e - e^* \in J(R)$. As in the preceding discussion, we can find a unique projection $f \in R$ such that $a - f \in U(R)$ such that $af = fa$, and therefore the result follows by Theorem 3.3. $\square$

We say that an ideal I of a $*$-ring R is a *$*$-ideal* provided that $I^* \subseteq I$. If I is a $*$-ideal of a $*$-ring, it is easy to check that R/I is also a $*$-ring.

Theorem 3.9 *Let I be a $*$-ideal of a $*$-ring R. If $I \subseteq J(R)$, then R is strongly J-$*$-clean if and only if*

(1) *R/I is strongly J-$*$-clean;*

(2) *R is abelian;*

(3) *Every idempotent lifts modulo I.*

Proof Suppose R is strongly J-$*$-clean. Then R is an abelian exchange ring, and so every idempotent lifts modulo I. For any $a \in R$, there exist a projection $e \in R$ and a unit $u \in R$ such that $a = e+u$; hence, $\overline{a} = \overline{e}+\overline{u}$ in R/I. Assume that there exist a projection $\overline{f} \in R/I$ and a unit $\overline{v} \in R/I$ such that $\overline{a} = \overline{f}+\overline{v}$. Then, we can find an idempotent $g \in R$ such that $f = g+r$ for some $r \in I$. Hence, $a = g + (v + r + t)$ for some $t \in J(R)$. Obviously, $g - g^* = f - r - f^* + r^* \in J(R)$. As $ag = ga, ag^* = g^*a$ and $v + r + t \in U(R)$, it follows from Lemma 3.8 that $g = e$, and so $\overline{f} = \overline{e}$ in R/I. Therefore R/I is strongly J-$*$-clean by Theorem 3.3.

Conversely, assume that $(1), (2)$ and (3) hold. For any $a \in R$, it follows from Theorem 3.3 that there exist a projection $\overline{e} \in R/I$ and a unit $\overline{u} \in R/I$ such that $\overline{a} = \overline{e} + \overline{u}$. As $e - e^2 \in I$, by hypothesis, there exists an idempotent $f \in R$ such that $e - f \in I$. Since every unit lifts modulo I, we may assume that $u \in U(R)$. Thus, $a = f + u + r$ for some $r \in I$. Set $v = u + r$. Then $a = f + v$ with $f = f^2 \in R, v \in U(R)$. As R is abelian, $af = fa$ and $af^* = f^*a$. Further, $f - f^* \equiv e - e^* \equiv 0 \pmod I$ and so $f - f^* \in J(R)$. Suppose that $a = g + w$ with $g = g^2 \in R, w \in U(R)$, $ag = ga, ag^* = g^*a$ and $g - g^* \in J(R)$. Then $\overline{a} = \overline{g} + \overline{w}$ in R/I. Clearly, R/I is uniquely clean, and so $f - g \in I \subseteq J(R)$. As $fg = gf$, we see that $(f - g)^3 = f - g$, and so $f = g$. In light of Lemma 3.8, R is strongly J-$*$-clean. $\square$

Let $P(R)$ be the intersection of all prime ideals of R, i.e., $P(R)$ is the *prime radical* of R. Recall that $a \in R$ is *strongly nilpotent* if for every sequence $a_0, a_1, \cdots, a_i, \cdots$ such that $a_0 = a$ and $a_{i+1} \in a_iRa_i$, there exists an n with $a_n = 0$. As is well known, the prime radical $P(R)$ is the set of all strongly nilpotent elements in R.

Corollary 3.10 *A $*$-ring R is strongly J-$*$-clean if and only if R is abelian and $R/P(R)$ is strongly J-$*$-clean.*

Proof Let $a \in P(R)$. For every sequence $a_0, a_1, \cdots, a_i, \cdots$ such that $a_0 = a^*$ and $a_{i+1} \in a_iRa_i$, we get a sequence $a_0^*, a_1^*, \cdots, a_i^*, \cdots$ such that $a_0^* = a$ and $a_{i+1}^* \in a_i^*Ra_i^*$. As $a \in R$ is strongly nilpotent, we can find some n such that $a_n^* = 0$, and so $a_n = 0$. This implies that a^* is strongly nilpotent; hence, $a^* \in P(R)$. We infer that $P(R)$ is a $*$-ideal. As every idempotent lifts modulo $P(R)$, we complete the proof by Theorem 3.9. $\square$

In [10], Li and Zhou proved that a $*$-ring R is strongly $*$-clean if and only if $R/J(R)$ is strongly $*$-clean, every projection is central and every projection lifts to a projection modulo $J(R)$. Analogous to the preceding discussion, we easily extend Li and Zhou's result as follows: A $*$-ring R is strongly $*$-clean if and only if $R/J(R)$ is strongly $*$-clean; R is abelian and every idempotent lifts modulo $J(R)$.

4 Extensions

By applying the preceding results, we will consider extensions of strongly J-$*$-clean rings. Let R be a $*$-ring, and let $R[i]=\{a+bi \mid a,b\in R, i^2=-1\}$. Then $R[i]$ is also a $*$-ring by defining $*: a+bi\mapsto a^*+b^*i$.

Lemma 4.1 *Let R be a ring with $2\in J(R)$. Then $U\big(R[i]\big)=\{a+bi \mid a+b\in U(R)\}$.*

Proof Assume that $(a+bi)(c+di)=1$. Then $ac-bd=1$ and $ad+bc=0$. Thus, $(a+b)(c+d)=ac+bd+ad+bc=(ac-bd)+2bd=1+2bd\in U(R)$. This implies that $a+b\in R$ is right invertible. As a result, we show that $U\big(R[i]\big)\subseteq\{a+bi \mid a+b\in U(R)\}$.

Assume that $a+b\in U(R)$. Then $a-b=(a+b)-2b\in U(R)$. Clearly, $a(a-b)^{-1}a=(a-b+b)(a-b)^{-1}(a-b+b)=\big(1+b(a-b)^{-1}\big)(a-b+b)=a+b+b(a-b)^{-1}b$. Therefore $a(a-b)^{-1}a(a+b)^{-1}-b(a-b)^{-1}b(a+b)^{-1}=1$. Likewise, $b(a-b)^{-1}a-a(a-b)^{-1}b=0$. It is easy to check that

$$\begin{aligned} & (a+bi)(a-b)^{-1}(a-bi)(a+b)^{-1} \\ = \;& \big(a(a-b)^{-1}a(a+b)^{-1}+b(a-b)^{-1}b(a+b)^{-1}\big)+\big(b(a-b)^{-1}a-a(a-b)^{-1}b\big)i \\ = \;& 1+2b(a-b)^{-1}b(a+b)^{-1} \\ \in \;& U(R). \end{aligned}$$

Thus, $a+bi\in R[i]$ is right invertible. Analogously, $(a-b)^{-1}(a-bi)(a+b)^{-1}(a+bi)\in U(R)$. Therefore $a+bi\in U\big(R[i]\big)$, as required. $\square$

Theorem 4.2 *Let R be a $*$-ring. Then $R[i]$ is strongly J-$*$-clean if and only if so is R.*

Proof Suppose that $R[i]$ is strongly J-$*$-clean. Then $2\in J(R)$. Further, every idempotent in $R[i]$ is a projection, and so is central. Let $a\in R$. Then we can find a projection $e+fi\in R[i]$ and a $u+vi\in J\big(R[i]\big)$ such that $a=(e+fi)+(u+vi)$ and $a(e+fi)=(e+fi)a$. Thus, $a=e+u$ and $ae=ea$. As $e+fi\in R[i]$ is an idempotent, $e\in R$ is central. Since $(e+fi)^*=e+fi$, we see that $e^*=e$. From $e+fi=(e+fi)^2$, we get $e^2-f^2=e$ and $2ef=f$. This implies that $(2e-1)f=0$, and then $f=0$. Hence, $e\in R$ is a projection. It is easy to verify that $u\in J(R)$, and therefore R is strongly J-$*$-clean.

Conversely, assume that R is strongly J-$*$-clean. Then R is an abelian exchange ring. In addition, every idempotent in R is a projection and $2\in J(R)$. Let $a+bi\in R[i]$. By hypothesis, there exist projections $e,f\in R$ and $u,v\in J(R)$ such that $a=e+u, b=f+v, ae=ea, bf=fb$. Thus, $a+bi=(e+f)+\big(u-f+(f+v)i\big)$. Clearly, $(e+f)^2-(e+f)=2ef\in J(R)$. As every idempotent lifts modulo $J(R)$, we can find an idempotent $g\in R$ such that $e+f=g+r$ where $r\in J(R)$. Thus, $a+bi=g+\big(r+u-f+(f+v)i\big)$ where $g=g^2=g^*$ and $(a+bi)g=g(a+bi)$.

Write $x = r + u - f$ and $y = f + v$. Then $x + y = r + u + v \in J(R)$. For any $c + di \in R[i]$, we see that $1 - (x + yi)(c + di) = (1 - xc + yd) - (xd + yc)i$. As $(1 - xc + yd) - (xd + yc) = (1 - xc - yd) - (xd + yc) + 2yd = 1 - (x + y)(c + d) + 2yd \in U(R)$. In light of Lemma 4.1, $1 - (x + yi)(c + di) \in U\big(R[i]\big)$, and so $x + yi \in J\big(R[i]\big)$. Therefore $R[i]$ is strongly J-$*$-clean, as asserted. □

Let R be a $*$-ring and G be a group. Then $*$ induces an involution of the group ring RG, denoted by $*$, where $(\sum_g a_g g)^* = \sum_g a_g^* g^{-1}$ (see [10, Lemma 2.12]). A group G is called *locally finite* if every finitely generated subgroup of G is finite. A group G is a 2-*group* if the order of each element of G is a power of 2.

Theorem 4.3 *Let R be a $*$-ring, and let G be a locally finite group. Then RG is a strongly J-$*$-clean ring if and only if R is a strongly J-$*$-clean ring and G is a 2-group.*

Proof Suppose that RG is strongly J-$*$-clean. Then it is uniquely clean. By virtue of [7, Theorem 12], R is uniquely clean and G is a 2-group. In view of Theorem 3.1, $2 \in J(RG)$, and so $2 \in J(R)$. In addition, every idempotent in RG is central, and so every idempotent in R is central in R. According to [7, Lemma 11], every idempotent in RG is in R. Therefore every idempotent in R is a projection. According to Theorem 3.3, R is strongly J-$*$-clean.

Conversely, assume that R is a strongly J-$*$-clean ring and G is a 2-group. By virtue of [7, Theorem 13], RG is uniquely clean. As R is strongly J-$*$-clean, we see that $2 \in J(R)$. Further, every idempotent in R is central. According to [7, Lemma 11], every idempotent in RG is in R. Therefore every idempotent in RG is a projection. By using Proposition 2.1, RG is strongly J-$*$-clean. □

Corollary 4.4 *Let R be a $*$-ring, and let G be a solvable group. Then RG is a strongly J-$*$-clean ring if and only if R is a strongly J-$*$-clean ring and G is a 2-group.*

Proof Suppose that RG is strongly J-$*$-clean. Then it is uniquely clean. By virtue of [7, Theorem 13], R is uniquely clean and G is a 2-group. As in the preceding discussion, we see that R is strongly J-$*$-clean.

Conversely, assume that R is a strongly J-$*$-clean ring and G is a 2-group. Analogously to the consideration in [7, Theorem 13], G is locally finite, and then the result follows from Theorem 4.3. □

Example 4.5 Let R be a Boolean ring and let G be a locally finite 2-group which is not abelian. Let $*$ be the identical automorphism of R. Then R is a $*$-ring and in view of Theorem 4.3 RG is a strongly J-$*$-clean ring which is not commutative.

References

[1] S.K. Berberian, Baer Rings and Baer $*$-Rings, U.S. Copyright Office, 1998.

[2] G. Borooah, A.J. Diesl and T.J. Dorsey, Strongly clean triangular matrix rings over local rings, J. Algebra, 312 (2007), 773-797.

[3] G. Borooah, A.J. Diesl and T.J. Dorsey, Strongly clean matrix rings over commutative local rings, J. Pure Appl. Algebra, 212 (2008), 281-296.

[4] H. Chen, On strongly J-clean rings, Comm. Algebra, 38 (2010), 3790-3804.

[5] H. Chen, On uniquely clean rings, Comm. Algebra, 39 (2011), 189-198.

[6] H. Chen, Rings Related Stable Range Conditions, Series in Algebra 11, World Scientific, Hackensack, NJ., 2011.

[7] J. Chen, W.K. Nicholson and Y. Zhou, Group rings in which every element is uniquely the sum of a unit and an idempotent, J. Algebra, 306 (2006) 453-460.

[8] J. Chen, Z. Wang and Y. Zhou, Rings in which elements are uniquely the sum of an idempotent and a unit that commute, J. Pure Appl. Algebra, 213(2009), 215-223.

[9] T.K. Lee and Y. Zhou, A class of exchange rings, Glasgow Math. J., 50 (2008), 509-522.

[10] C. Li and Y. Zhou, On strongly $*$-clean rings, J. Algebra Appl., 10 (2011), 1363-1370.

[11] W.K. Nicholson, Clean rings: a survey, Advances in Ring Theory, World Sci. Publ., Hackensack, NJ, 2005, 181-198.

[12] W.K. Nicholson and Y. Zhou, Rings in which elements are uniquely the sum of an idempotent and a unit, Glasgow Math. J., 46 (2004), 227-236.

[13] J. Stock, On rings whose projective modules have the exchange property, J. Algebra, 103 (1986), 437-453.

[14] L. Vaš, $*$-Clean rings; some clean and almost clean Baer $*$-rings and von Neumann algebras, J. Algebra, 324 (2010), 3388-3400.

[15] X. Wang, Uniquely strongly clean group rings, Comm. Math. Research, 28 (2012), 17-25.

[16] K.N. Rajeswari and R. Aziz, Certain units in $M_2(\mathbb{Z})$ which are uniquely clean, Research J. Pure Algebra, 1 (2011), 58-70.

[17] X. Yang, A survey of strongly clean rings, Acta Appl. Math., 108 (2009), 157-173.